

Rec'd PCT/PTO 13 DEC 2004

PCT/EP 03/50276 #2

10/517788 9. 2003

PA 1017988

RECD 02 OCT 2003

PTO

PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

June 04, 2003

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/393,149

FILING DATE: July 02, 2002

PRIORITY  
DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS

  
*P. R. Grant*

P. R. GRANT  
Certifying Officer

BEST AVAILABLE COPY

07/02/02  
1074 U.S. PTO

7-03-02 1074 U.S. PTO 1074 U.S. PTO / 1074 U.S. PTO

1074 U.S. PTO  
1074 U.S. PTO

## PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53 (c).

		DOCKET NUMBER	JAB-1712
INVENTOR(S) / APPLICANT(S)			
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Van Dun Wouters Janicot Mocek Laakso	Jacobus Walter Michel Ursula Jodi	A.J. B.L. M.F. M. A.	Kasterlee, Belgium Kapellen, Belgium Mol, Belgium Bellevue, Washington, USA North Bend, Washington, USA
TITLE OF THE INVENTION (280 characters max)			
A NOVEL BAFILEOMYCIN-LIKE METABOLITE FROM A NOVEL Micromonospora SPECIES			
CORRESPONDENCE ADDRESS			
Direct all correspondence to: <input checked="" type="checkbox"/> Customer Number 000027777 OR <input type="checkbox"/> Firm of Individual Name:			
ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> Specification	Number of Pages	20	<input type="checkbox"/> Application Data Sheet
<input checked="" type="checkbox"/> Claims	Number of Claims	10	<input type="checkbox"/> CD(s), Number
<input checked="" type="checkbox"/> Drawing(s)	Number of Sheets	4	<input type="checkbox"/> Other (specify) <input type="text"/>
METHOD OF PAYMENT (check one)			
<input type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees.		Provisional Filing Fee Amount (\$)	\$ 160.00
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit any overpayment to Deposit Account No. 10-0750			

The Invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

No

Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

SIGNATURE: Mary A. Appolina

REGISTRATION NO. 34,087

TYPED or PRINTED NAME Mary A. Appolina

DATE: July 2, 2002

TELEPHONE (732) 524-3742

**PROVISIONAL APPLICATION FILING ONLY**

DOCKET NO. JAB-1712

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: VAN DUN et al

For : A NOVEL BAFILOMYCIN-LIKE METABOLITE FROM A NOVEL  
Micromonospora SPECIES

Express Mail Certificate

"Express Mail" mailing number: EF195553720US

Date of Deposit: July 2, 2002

I hereby certify that this complete application, including specification pages, claims and informal drawings, is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Commissioner for Patents, Washington, D.C. 20231.

Judy Ubry

(Typed or printed name of person mailing paper or fee)

Judy Ubry  
(Signature of person mailing paper or fee)

A NOVEL BAFILOMYCIN-LIKE METABOLITE FROM A NOVEL  
*Micromonospora* SPECIES


5 The present invention relates to a novel baflomycin-like metabolite isolated from a novel *micromonospora* species having potent antiproliferative activity as well as to processes for isolating said metabolite, pharmaceutical compositions containing this metabolite and methods of treatment using said metabolite.

10 Although the increasing basic knowledge of major cellular signaling pathways has allowed recent discovery of promising "targeted therapeutics" to help fighting cancer (e.g., inhibitors of farnesyltransferase, growth factor receptor tyrosine kinases, Bcr/abl, etc...), more classical cytotoxics still remain an important part of the foundation of an oncology product portfolio in the modern pharmaceutical industry. The partial success 15 of traditional cancer cytotoxic chemotherapy stems in large part from agents derived from natural sources.

Macrolides are a group of antibiotics that includes natural members. Most of the macrolides are produced by *Streptomyces* spp. bacteria. However also macrolides 20 produced by *Micromonospora* spp. are known. Macrolides are characterised in that they are composed of a large aglycone ring (from 14-16 carbon atoms) to which are attached several sugars. A special series of these macrolides are the baflomycins. These compounds share the structural feature of having a 16-membered macrocyclic lactone ring and one or two side-chain sugars. The aforementioned baflomycins were 25 previously known to have activity as antiparasitics, antithelmintics, antibiotics, insecticides, antifungals, agents for the therapy of bone diseases and as vacuolar-type ATPase proton pump (v-ATPase) inhibitors. Later on, it has been found that, when run through whole tumour-cell assays, compounds from this group exhibit antitumour activity.

30

WO 01/02413 discloses baflomycin derivatives of formula (II),



wherein R is -H or -CO-CH=CH-COOH, and pharmaceutically acceptable salts and esters thereof. The compounds display activity against tumour growth in mammals.


WO 01/02413 further provides a method for the treatment or prophylaxis of tumour

5 invasion by administration of compounds of formula (II) or of known macrolide compounds.

The present invention encompasses a novel structure of formula (I) or a

pharmaceutically acceptable salt thereof, with a new hexadiene sidechain at C23. The compound of formula (I) shows antiproliferative activity and is most unlikely a substrate for P-glycoprotein and/or Multidrug Resistant Protein.

The compound of formula (I) is a new baflomycin-like derivative isolated from liquid fermentation cultures of novel actinomycetes of the genus *Micromonospora*.



15

The compound of formula (I) was extensively characterized for its ability to inhibit tumour cell growth. The dose-dependent inhibition of the compound of formula (I) in four selected tumour cell proliferation assays was typical for classical cytotoxics, with

-3-

very steep and narrow dose-response curves. Initial observation on inhibition of tumour cell proliferation was extended to a wide variety of human tumour cell lines, from ovary, prostate, pancreas, colon and lung origins, with observed IC<sub>50</sub> values in the nM range of concentrations.

5 Multi-drug resistance is considered as one of the major elements in the failure of anticancer therapeutics in clinical trials. Patients become resistant to therapeutic agents by overexpression of both P-glycoprotein and Multidrug Resistant Protein which represent the major elements of the Multidrug Resistant system. In addition,

10 expression of such proteins in the intestinal epithelium markedly reduces compound absorption in the case of oral administration of the drug. The activity of the compound of formula (I) on P-glycoprotein or Multidrug Resistant Protein-expressing tumour cell lines and respective parental cells, was compared to that of a known substrate for P-glycoprotein/Multidrug Resistant Protein, adriamycin. The compound of formula (I)

15 inhibited cell proliferation in the nM range of concentrations regardless of the P-glycoprotein and/or Multidrug Resistant Protein expression status. These results strongly suggest that there is no substantial recognition of the compound of formula (I) by the P-glycoprotein and Multidrug Resistant Protein pump proteins.

20 The present novel cancer therapeutic agent shows, like most members of this class of compounds, some acute toxicity at higher concentrations. Persons skilled in the art will however be able to determine an appropriate therapeutic range depending upon the condition of the patient.

25 Pharmaceutically acceptable addition salts encompass pharmaceutically acceptable acid addition salts and pharmaceutically acceptable base addition salts. The compound of formula (I) having basic properties may be converted in its pharmaceutically acceptable acid addition salts by treating it with an appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic

30 acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-amino-salicylic, pamoic and the like acids.

35 The compound of formula (I) having acidic properties may be converted in its pharmaceutically acceptable base addition salt by treating it with a suitable organic or inorganic base. Appropriate base salt forms comprise, for example, the ammonium

-4-

salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-*D*-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

5

The term "acid or base addition salts" also comprises the hydrates and the solvent addition forms which the compound of formula (I) is able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

10 This invention also provides a method for inhibiting tumour growth, a method for the treatment of cancer, by administering an effective amount of the compound of the present invention, to a subject, e.g. a mammal (and more particularly a human) in need of such treatment. Examples of cancer types which may be inhibited, but are not limited to solid and haematological cancers, leukemia, skin cancers such as melanoma and  
15 basal cell carcinoma, renal cancer, lung cancer such as small cell lung cancer, non-small-cell lung cancer and alveolar cell carcinoma, colon cancer, breast cancer, prostate cancer, pancreatic cancer, stomach cancer and medullary thyroid cancer.

The present compound, in view of its antibacterial characteristics, is also useful as an antibacterial medicine. This invention also provides a method for treating bacterial  
20 infections, by administering an effective amount of the compound of the present invention, to a subject, e.g. a mammal (and more particularly a human) in need of such treatment.

25 The present invention discloses the compound of formula (I) for use as a medicine as well as the use of these compounds of formula (I) for the manufacture of a medicament for inhibiting tumour growth.

In view of its useful pharmacological properties, the subject compound may be formulated into various pharmaceutical forms for administration purposes.

30

To prepare the pharmaceutical compositions of this invention, an effective amount of a particular compound, in base or acid addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired  
35 for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form,

-5-

any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, 5 pills, capsules and tablets.

Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise 10 sterile water, at least in large part, though other ingredients, to aid solubility for example, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the 15 compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause a significant deleterious effect to the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired 20 compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on or as an ointment.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. 25 Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient, calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, 30 injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

Those skilled in the art could easily determine the effective amount from the test results presented hereinafter. In general it is contemplated that a therapeutically effective 35 amount would be from 0.001 mg/kg to 100 mg/kg body weight, and in particular from 0.5 mg/kg to 100 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day.

Said sub-doses may be formulated as unit dosage forms, for example, containing 0.5 to 500 mg, and in particular 10 mg to 500 mg of active ingredient per unit dosage form.

Brief description of the drawings

**FIG 1A:** *Dose-dependent inhibition of tumour cell proliferation (bioassay) by the compound of formula (I).* Human MCF-7 tumour cells were treated with indicated concentrations of the compound of formula (I) for 4 days. Results, expressed as % of control untreated cells, are the mean  $\pm$  SE of at least 3 different experiments.

**FIG 1B:** *Dose-dependent inhibition of tumour cell proliferation (bioassay) by the compound of formula (I).* Human HT-29 tumour cells were treated with indicated concentrations of the compound of formula (I) for 4 days. Results, expressed as % of control untreated cells, are the mean  $\pm$  SE of at least 3 different experiments.

**FIG 1C:** *Dose-dependent inhibition of tumour cell proliferation (bioassay) by the compound of formula (I).* Human K562/C1,000 tumour cells were treated with indicated concentrations of the compound of formula (I) for 4 days. Results, expressed as % of control untreated cells, are the mean  $\pm$  SE of at least 3 different experiments.

**FIG 1D:** *Dose-dependent inhibition of tumour cell proliferation (bioassay) by the compound of formula (I).* Human Malme-3M tumour cells were treated with indicated concentrations of the compound of formula (I) for 4 days. Results, expressed as % of control untreated cells, are the mean  $\pm$  SE of at least 3 different experiments.

20 Experimental Part

In the course of a screening program designed to identify new antitumour compounds (cytotoxics), a new baflomycin-like metabolite was isolated. The same metabolite was isolated from liquid fermentation cultures of two different actinomycete of the genus *Micromonospora*. *Micromonospora spp.* JS1035 was isolated from river sediments from Sombo, Cameroon while *Micromonospora spp.* JS1044 was isolated from river sediments from Mundemba, Cameroon. As soon as it was detected that both strains produced the same active compound it was decided to further characterize the compound present in liquid fermentation cultures from *Micromonospora spp.* JS1035.

30 A culture of each micro-organism utilised for the production of these new macrolide have been deposited in the Laboratory for Microbiology in the Belgian Coordinated Collections of Microorganism at the University of Gent under accession number LMG P-21525 for *Micromonospora spp.* JS1035 and under accession number LMG P-21526 for *Micromonospora spp.* JS1044. These deposits have been made under the provisions

-8-

of the Budapest Treaty and all restrictions on the availability thereof to the public will be irrevocably maintained upon granting of a patent on this application.

General

5

Fermentation and Isolation

Liquid fermentation cultures of *Micromonospora spp.* JS1035 were extracted with ethyl acetate to yield the crude extract that was subsequently purified using counter current partition chromatography (CPC). Pure material for structure elucidation was obtained 10 after two consecutive CPC fractionations using different conditions. Mass spectral analysis (see Table 1) of this material indicated a molecular weight of 853 and a molecular formula of C<sub>47</sub>H<sub>67</sub>NO<sub>13</sub>.

15

**Table 1.** *Physico-chemical properties of the compound of formula (I).*

Appearance	Light brown solid
Melting Point	114-120°C
[ $\alpha$ ] <sup>25</sup> <sub>D</sub>	-38.0 (c 0.1, MeOH)
Molecular Formula	C <sub>47</sub> H <sub>67</sub> NO <sub>13</sub>
Pos. ESI*-MS	854 (M+H) <sup>+</sup>
HRFAB*-MS (m/z)	Found: 876.4514 Calcd: 876.4510 (for C <sub>47</sub> H <sub>67</sub> NO <sub>13</sub> Na)
UV (MeOH) $\lambda_{\text{max}}$ nm ( $\epsilon$ )	233, 286, 345
IR* $\nu_{\text{max}}$ (film) cm <sup>-1</sup>	3408, 2923, 1717, 1687, 1614, 1247, 1097, 751

\* ESI: Electron Spray Ionisation, HRFAB: High Resolution Fast Atom Bombardment, IR: Inversion Recovery

Structure Determination

20

The NMR data for the compound of formula (I) was initially obtained in CDCl<sub>3</sub>. However, it was soon discovered that the compound of formula (I) was not stable in this solvent. Subsequent NMR work was performed using acetone-d<sub>6</sub> and yielded a complex proton NMR spectrum. A full set of NMR experiments including <sup>13</sup>C NMR, Distortionless Enhancement by Polarization Transfer (DEPT), Correlated Spectroscopy (COSY), Heteronuclear Multiple Quantum Coherence (HMQC), and Heteronuclear Multiple Band Coherence (HMBC) was obtained.

25

Assembling the partial structures for the compound of formula (I) was straightforward. The NMR data (Table 2) for the macrolide, C16-C23, and C1'-C4' were in good

agreement with published data. (Werner G. *et al.* J. Antibiotics 37: 110-117, 1984) Attachment of the new hexadiene sidechain (C24-C29) was accomplished by careful study of the correlations between relevant atoms. For instance, H24 showed an HMBC correlation to C22 and C23, and H25 also showed an HMBC correlation to C23. The 5 COSY data were also consistent with placement of the new sidechain at C23. Several  $^{13}\text{C}$  NMR signals (C5', C6', and C9') corresponding to the cyclopentenone ring were obscured in the  $^{13}\text{C}$  NMR spectrum, probably due to the presence of resonance structures between the vinylic OH and the carbonyl group.

10 Other published reports detail the stereochemistry for other bafilomycins and other members of this class (O'Shea M. *et al* J. Antibiotics 50: 1073-1077, 1997). Molecules in this class have the same stereochemistry for the lactone ring, the tetrahydropyrin ring, and the C16-C18 linkage. All appear to possess a hydrogen-bonding network that includes C19OH, C17OH, and C1=O. This network effectively defines the 15 configuration of the side chain. Based on the analysis of coupling constant, chemical shift, and Rotating Frame Overhauser Effect Spectroscopy (ROESY) data, the proposed stereochemistry of the compound of formula (I) is consistent with this conformation. The new hexadiene sidechain at C23 was assigned as alpha to the ring based on two key ROESY correlations. There is a correlation between H23 and CH<sub>3</sub>-36 and also 20 between H23 and H21. These correlations would be unlikely unless H23 was positioned above the ring.

**Table 2.**  $^1\text{H}$  and  $^{13}\text{C}$  NMR chemical shifts for the compound of formula (I)<sup>a</sup>

	$\delta_{\text{C}}$ (ppm)	M	$\delta_{\text{H}}$ (ppm)
1	167.57	s	
2	141.97	s	
3	133.35	d	6.63 (1H, s)
4	132.74	s	
5	145.55	d	5.94 (1H, d, 8.7)
6	37.85	d	2.52 (1H, m)
7	80.17	d	3.28 (1H, br d, <i>J</i> 5.4)
8	41.93	d	1.86 (1H, m)
9	42.27	t	2.12 (1H, m), 2.03 (1H, m)
10	144.98	s	
11	125.19	d	5.80 (1H, d, 10.7)
12	134.53	d	6.67 (1H, dd, 15.0, 10.8)
13	126.95	d	5.12 (1H, m)
14	82.98	d	4.07 (1H, m)

-10-

15	77.01	d	4.96 (1H, dd, 8.4, 1.3)
16	38.23	d	2.13 (1H, m)
17	70.80	d	4.11 (1H, m)
18	42.27	d	1.84 (1H, m)
19	100.26	s	
20	40.22	t	2.35 (1H, dd, 11.7, 4.8), 1.34 (1H, m)
21	75.11	d	5.08 (1H, m)
22	41.81	d	1.50 (1H, m)
23	75.34	d	4.14 (1H, m)
24	130.81	d	5.50 (1H, dd, 15.1, 7.6)
25	133.51	d	6.18 (1H, dd, 15.5, 15.4)
26	129.72	d	5.99 (1H, dd, 15.1, 10.5)
27	137.21	d	5.70 (1H, m)
28	26.16	t	2.09 (2H, m)
29	13.73	q	0.97 (3H, d, 2.8)
30	14.13	q	1.94 (3H, br s)
31	17.62	q	1.03 (3H, d, 7.0)
32	22.20	q	0.90 (3H, d, 6.8)
33	20.33	q	1.93 (3H, br s)
34	10.02	q	0.85 (3H, m)
35	7.2	q	1.00 (3H, d, 2.5)
36	11.61	q	0.85 (3H, m)
2-OCH <sub>3</sub>	59.94	q	3.50 (3H, s)
14-OCH <sub>3</sub>	55.63	q	3.23 (3H, s)
1'	168.14	s	
2'	134.97	d	7.57 (1H, d, 15.3)
3'	132.97	d	6.82 (1H, d, 15.3)
4'	165.00	s	
5'	b	s	
6'	b	S	
7'	27.75	t	2.52 (2H, m)
8'	37.31	t	2.52 (2H, m)
9'	b	s	

<sup>a</sup>δ in ppm, J value in Hz. Data were recorded at 500 and 125 MHz, respectively in acetone-d<sub>6</sub>. <sup>b</sup>Signal not observed.

#### Biological Activity

5 A panel of cell proliferation assays (referred to herein as "bioassays") based on the inhibition of cell growth in selected human tumour cell lines (mammary MCF-7, colon HT-29, leukemia K562/C1,000 and melanoma Malme-3M) was used to identify antiproliferative activities in the different samples throughout the purification

-11-

procedures. Extracts from strain JS1035 grown in four different media were tested in bioassays; the one that contained most of the antiproliferative activity was selected for further study. A 1:1,000-dilution of the active extract resulted in an almost complete inhibition of cell proliferation in the three cell lines tested (melanoma Malme-3M cell line was omitted at this stage). From this initial crude extract, collected fractions from counter current partition chromatography were individually analyzed on MCF-7 cell proliferation. A major single fraction was shown to contain antiproliferative activity, which was then further purified by HPLC. A single HPLC peak was found to contain antiproliferative activities as measured by inhibition of cell proliferation in the bioassay. This peak was used to initiate structure elucidation of the active compound of formula (I). Subsequent structure work was completed using material isolated with two consecutive CPC steps.

The purified compound of formula (I) was extensively characterized for its ability to inhibit tumour cell growth. As indicated in *Figure 1A-D*, the dose-dependent inhibition of the compound of formula (I) in the four selected tumour cell proliferation assays was typical for classical cytotoxics, with very steep and narrow dose-response curves. The compound of formula (I) exhibited similar potency for inhibition of tumour cell proliferation in the nM range of concentrations, with a slightly higher activity on colon HT-29 carcinoma cells (sub-nanomolar concentrations) (*Table 3*). Initial observation on inhibition of tumour cell proliferation was extended to a wide variety of human tumour cell lines, from ovary, prostate, pancreas, colon and lung origins, with observed IC<sub>50</sub> values in the nM range of concentrations (*Table 4*).

The activity of the compound of formula (I) on P-glycoprotein- or Multidrug Resistant Protein-expressing tumour cell lines and respective parental cells, K562/C1,000+ and K562A7, COR-L23/R and COR-L23, was compared to that of a known substrate for P-glycoprotein/Multidrug Resistant Protein, adriamycin. As indicated in *Table 5*, the compound of formula (I) inhibited cell proliferation in the nM range of concentrations regardless of the P-glycoprotein and/or Multidrug Resistant Protein expression status (no significant difference between IC<sub>50</sub> values). On the contrary, K562/C1,000+ and CORL23/R cells exhibited a 250-fold and 20-fold resistance to cell killing by adriamycin, respectively, as compared to their respective parental cell lines. These results strongly suggested that there is no substantial recognition of the compound of formula (I) by the P-glycoprotein and Multidrug Resistant Protein pump proteins.

-12-

**Table 3.** *Inhibition of tumour cell proliferation by the compound of formula (I).*

Sensitivity to cell killing by the compound of formula (I) was assessed in breast MCF-7, colon HT-29, leukemia K562/C1,000 and melanoma Malme-3M cells. Cell viability was determined after a 4-day incubation in the absence (control) or presence 5 of increasing concentrations of the compound of formula (I). Results, means  $\pm$  SE from at least three separate experiments, are expressed as IC<sub>50</sub> values (in nM).

<i>Cell Lines</i>	<i>Compound of formula (I)</i>
<b>MCF-7</b>	$1.92 \pm 0.35$
<b>HT-29</b>	$0.61 \pm 0.04$
<b>K562/C1,000</b>	$9.89 \pm 7.35$
<b>Malme-3M</b>	$7.96 \pm 2.73$

**Table 4.** *Effect of the compound of formula (I) on human tumour cell proliferation.*

10 Activity of the compound of formula (I) on tumour cell proliferation was assessed on several human tumour cell lines from various origins. As indicated in the *Materials and Methods* section, a 4-day incubation period in the absence (control) or presence of increasing concentrations of the compound of formula (I) was followed by cell viability determination using MTT staining method. Results, means  $\pm$  SE from at least three 15 separate experiments, are expressed as IC<sub>50</sub> values (in nM) which represent the concentration of compound required to inhibit cell proliferation by 50%.

<i>Cell Line</i>	<i>Tissue origin</i>	<i>Compound of formula (I)</i>
<b>A2780</b>	ovary	$1.37 \pm 0.13$
<b>DU145</b>	prostate	$1.95 \pm 0.30$
<b>CAPAN-1</b>	pancreas	$6.93 \pm 3.39$
<b>HCT116</b>	colon	$1.56 \pm 0.03$
<b>LoVo</b>	colon	$5.98 \pm 5.52$
<b>H1299</b>	lung	$1.46 \pm 0.13$

20 **Table 5.** *Is the compound of formula (I) a substrate for P-glycoprotein/Multidrug Resistant Protein?* Sensitivity to cell killing by the compound of formula (I) and adriamycin was assessed in human K562A7 parental leukemia cells and P-glycoprotein overexpressing counterpart, K562/C1,000+ cells; as well as parental small cell carcinoma CORL23 cells, and Multidrug Resistant Protein overexpressing counterpart,

-13-

CORL23/R cells. Cell viability was determined after a 4-day incubation in the absence (control) or presence of increasing concentrations of the compound of formula (I). Results, mean  $\pm$  SE from at least two separate experiments, are expressed as IC<sub>50</sub> values (in nM).

5

<i>Cell Line</i>	<i>Compound of formula (I)</i>	<i>Adriamycin</i>
<b>K562A7</b>	$14.10 \pm 1.25$	$26.78 \pm 5.82$
<b>K562/C1,000+</b>	$29.34 \pm 1.80$	$7,829 \pm 806$
<b>COR-L23</b>	$5.13 \pm 0.91$	$36.57 \pm 19.09$
<b>COR-L23/R</b>	$6.90 \pm 0.24$	$676 \pm 261$

### Example 1

#### Structure Determination

10 <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded at 300K on a Bruker Avance DRX 500 spectrometer operating at 500 and 125 MHz, respectively. Standard pulse sequences were used for DEPT, COSY, HMQC, and HMBC experiments. ROESY data was acquired using a spinlock pulse of 300,000  $\mu$ s (at 22 dB). Melting points were determined using an Electrothermal 9100 melting point apparatus and are uncorrected.

15 Inversion Recovery (IR) data was acquired using a Perkin Elmer 1600 Series FTIR spectrophotometer. A Perkin-Elmer 243B polarimeter was used to procure optical rotation data. High resolution MS data was obtained using a Micromass 70SEQ Tandem Hybrid Mass Spectrometer. CPC was performed on a P.C., Inc. high speed countercurrent chromatograph, equipped with an Ito multilayer coil column interfaced

20 with a Waters 991 Photodiode Array detector (monitoring at 270 nm).

#### Fermentation and Isolation

The actinomycete strain JS1035 was isolated from river bottom sediment collected near Sombo, Cameroon. The seed medium contained glucose 20 g, Pharmamedia (Trader's Protein) 15.0 g, yeast extract 5.0 g, CaCO<sub>3</sub> 4.0 g, (NH<sub>4</sub>)<sub>2</sub>(SO<sub>4</sub>) 3.0 g, ZnSO<sub>4</sub>·7H<sub>2</sub>O 0.03 g in 1 L distilled water, pH adjusted to 6.5. The production medium contained glucose

-14-

20.0 g, dextrin 50.0 g, Pharmamedia (Trader's Protein) 30.0, yeast extract 1.0 g, CaCO<sub>3</sub> 5.0 g, CoCl<sub>2</sub>·6H<sub>2</sub>O 0.001 g in 1 L distilled water, pH adjusted to 7.0. The media were dispensed in 25 or 30 mL aliquots to 250 mL flasks and autoclaved for 15 min. Frozen spore stocks (1.0 mL) were used to inoculate starter cultures (25 mL seed media per 5 250 mL Erlenmeyer flask). Starter cultures were incubated at 28°C and ~ 75% humidity on an orbital shaker (2 " throw, 250 rpm) for 48 h. The starter cultures (1 mL) were used to inoculate 200 production media flasks (30 mL production media per 250 mL Erlenmeyer flask), which were then incubated for 6 days at the above conditions.

10 10 Ethyl acetate (22.5 mL) was added to each flask, and the flask contents were pooled into 500 mL centrifuge bottles. Each bottle was shaken vigorously and centrifuged at 6000 x g for 8 minutes. The ethyl acetate was removed from each bottle and dried over anhydr. Na<sub>2</sub>SO<sub>4</sub>. Ethyl acetate was again added to the centrifuge bottles to a total volume of approximately 450 mL, and the process was repeated. The extract was dried 15 under reduced pressure to yield a crude extract (5.4 g).

An aliquot (400 mg) of the crude extract was subjected to CPC. The solvent system consisted of an equilibrated mixture of n-hexane, EtOAc, MeOH, and water (1:3:3:3, v/v/v/v). The lower phase was used as the stationary phase, and the upper phase served as the mobile phase which was pumped at 3 mL/minute. The column rotation speed 20 was 800 rpm. The crude extract was prepared by dissolving it in a mixture of upper and lower phases. The material was then loaded onto the column. The first peak to elute was collected and dried to yield 303 mg of purified material. This purified material was again subjected to CPC chromatography using modified conditions. The modified solvent system consisted of an equilibrated mixture of water, MeOH, and n-hexane (1:9:10, v/v/v). Again, the lower phase was used as the stationary phase, and 25 the upper phase served as the mobile phase. After 90 minutes, the upper and lower phases were switched and the first peak to elute after the solvent change was collected and dried to yield 9.2 mg of the pure compound of formula (I).

Initially, the compound of formula (I) was isolated using HPLC following the first CPC 30 separation. In that case, the active CPC fraction was further separated using reversed-phase HPLC (LiChrospher, 10 µm, 10 x 25 mm, C<sub>18</sub>). A linear gradient starting at 95:5 CH<sub>3</sub>CN:H<sub>2</sub>O to 100% CH<sub>3</sub>CN over 25 minutes was used to isolate the compound of

-15-

formula (I). A 5-mg HPLC injection yielded 0.5 mg of compound (I).

### Drugs

All samples tested were dissolved in DMSO and further dilutions (typically 1:1,000) 5 were made in culture medium, with final DMSO concentrations never exceeding 0.1 % (v/v) in cell proliferation assays. MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide], MTS [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium inner salt] and PMS [phenazine methosulfate] were dissolved in phosphate-buffered saline (PBS).

10

### Cell lines and cell culture

The human MCF-7 mammary adenocarcinoma cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 2 mM L-glutamine, 1 mM NaPyruvate, 50 µg/ml gentamicin and 5 % fetal calf serum. The human HT-29 colon 15 carcinoma cells were cultured in Mc Coy's 5a medium supplemented with 2 mM L-glutamine, 50 µg/ml gentamicin and 5 % fetal calf serum. The human Malme-3M melanoma cells were cultured in DMEM medium supplemented with 2 mM L-glutamine, 1 mM NaPyruvate, 50 µg/ml gentamicin and 10 % fetal calf serum. The human DU 145 prostate carcinoma cells were cultured in DMEM medium 20 supplemented with 2 mM L-glutamine, 1 mM NaPyruvate, 50 µg/ml gentamicin and 10 % fetal calf serum. The human Capan-1 pancreatic adenocarcinoma cells were cultured in DMEM medium supplemented with 2 mM L-glutamine, 1 mM NaPyruvate, 50 µg/ml gentamicin and 10 % heat inactivated fetal calf serum. The human HCT 116 25 colorectal carcinoma cells were cultured in Mc Coy's 5a medium supplemented with 2 mM L-glutamine, 50 µg/ml gentamicin and 10 % fetal calf serum. The human H1299 non-small lung carcinoma cells were cultured in RPMI 1640 medium supplemented with 2 mM L-glutamine, 1 mM NaPyruvate, 10 mM HEPES and 10 % heat inactivated fetal calf serum. All three cell lines were obtained from ATCC [American Type Culture Collection, Manassas, VA, USA]. The human A2780 ovarian carcinoma cell 30 line was a kind gift from Dr. T.C. Hamilton [Fox Chase Cancer Centre, Pennsylvania, USA] and was cultured in RPMI 1640 medium supplemented with 2 mM L-glutamine, 50 µg/ml gentamicin and 10 % fetal calf serum. The human LoVo colon

-16-

adenocarcinoma cell line was a kind gift from Dr. M. Grandi [Pharmacia-Fice, Nerviano, Milano, Italy] and was cultured in HAM's F12 medium supplemented with 2 mM L-glutamine, 50 µg/ml gentamicin, 1 % Vitamins (BME) and 10 % fetal calf serum. Cells were routinely kept as monolayer cultures at 37°C in a humidified 5 % CO<sub>2</sub> atmosphere. The human K562/C1,000 leukemia cell line was a kind gift by Dr. H. Heyligen [Dr. Willems Instituut, Diepenbeek, Belgium] and is a P-glycoprotein expressing cell line obtained by culturing the cells at increasing concentrations of colchicine. The cells were kept as a suspension culture at 37°C in a humidified 5 % CO<sub>2</sub> atmosphere in RPMI 1640 medium supplemented with 2 mM L-glutamine, 50 µg/ml gentamicin and 5 % fetal calf serum. The human K562A7 leukemia cell line was a kind gift by Dr. H. Heyligen and is the parent, drug sensitive cell line of the K562/C1,000 cells. The K562/C1,000+ cell line was obtained by culturing the K562/C1,000 cells for an additional 2 weeks with 1 µM colchicine. This treatment induces higher levels of P-glycoprotein and mRNA and a more stable resistance profile. The cells were kept as a suspension culture at 37°C in a humidified 5 % CO<sub>2</sub> atmosphere in RPMI 1640 medium supplemented with 2 mM L-glutamine, 50 µg/ml gentamicin and 5 % fetal calf serum. The human large cell lung cancer cell lines were obtained from ECACC (European Collection of Cell Cultures, Salisbury, U.K.). COR-L23 is the parent, drug sensitive cell line and COR-L23/R is the Multidrug Resistant Protein-expressing Multidrug Resistant-variant, developed by continuous step-wise *in vitro* incubation of COR-L23 cells with increasing concentrations of doxorubicin. The culture medium for both cell lines consisted of RPMI 1640 supplemented with 2 mM L-glutamine, 50 µg/ml gentamicin and 10% fetal calf serum. Cells were routinely kept as monolayer cultures at 37°C in a humidified 5% CO<sub>2</sub> atmosphere.

All media and supplements were obtained from Life Technologies, Merelbeke, Belgium. Cells were free of mycoplasma contamination as determined using the Gen-Probe *Mycoplasma* Tissue Culture kit [BioMérieux, Brussel, Belgium].

-17-

Cell proliferation assays


Cell proliferation assays were performed as described (A critical assessment of the use of microculture Tetrazolium assays to measure cell growth and function (Mossman, T. J. Immunol Meth 65: 55-63, (1983)).

- 5 In case of adherent cell lines, cells were seeded in Falcon/NUNC 96-well culture plates [Life Technologies, Merelbeke, Belgium] and allowed to adhere to the plastic for 18-48 hr. Medium was then changed and drugs and/or solvents were added. Following 4-day incubation, cell density was assessed using a MTT-based assay measuring absorbance at 540 nm using a Emax 96-well spectrophotometer [Sopachem, Brussel, Belgium].
- 10 For the experiments with the suspension culture, K562A7, K562/C1,000 and K562/C1,000+ cells were seeded in Falcon® 96-well culture plates. Test drugs and/or medium were added immediately after seeding. Following 4-day incubation, cell growth was assessed using a MTS based assay measuring absorbance at 490 nm using an Emax 96-well spectrophotometer.
- 15 The antiproliferative activity is calculated as percentage of remaining viable cells in treated *versus* control (untreated cells) conditions. Results are expressed as IC<sub>50</sub> values which represent the concentration of compound required to inhibit cell proliferation by 50%. Within an experiment, the result for each experimental condition is the mean of 3 replicate wells.

-18-

## CLAIMS

## 1. A compound of formula (I)



5 or a pharmaceutically acceptable salt thereof

10 2 A pharmaceutical composition comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as described in claim 1.

15 3 A process for preparing a pharmaceutical composition as claimed in claim 2 wherein a therapeutically effective amount of a compound as claimed in claim 1 is intimately mixed with a pharmaceutically acceptable carrier.

20 4 A process for the production of a compound of formula (I) as defined in claim 1, comprising cultivating a strain of a microorganism capable of producing a compound of formula (I), recovering the compound of formula (I) from the cultured broth, and optionally, salifying the recovered compound.


25 5 A process according to claim 4, wherein the microorganism is a *Micromonospora* species.

6 A compound according to claim 1 for use as a medicine.

25 7 The use of a compound according to claim 1 in the manufacture of a medicament for inhibiting cancer or bacterial growth.

-19-


- 8 A method for the treatment of cancer, comprising administering to a mammal in need of such treatment an effective amount of a compound of formula (I) as defined in claim 1.
- 5
9. A compound isolated from *Micromonospora* spp JS1035 deposit No LMG P-12525 having antitumour activity.
10. A compound as defined in claim 9 wherein the compound has a molecular weight of 853 and a molecular formula of  $C_{47}H_{67}NO_{13}$ .

ABSTRACT5 A NEW BAFILOMYCIN-LIKE METABOLITE FROM A NOVEL  
*Micromonospora* SPECIES

10 The invention provides a compound of formula (I) and pharmaceutically acceptable salts thereof. Processes for the isolating of this bafilomycin metabolite from a novel *Micromonospora* species, pharmaceutical compositions containing this metabolite and methods of treatment using said metabolite are also described


-1/4-

FIG 1A




-2/4-

FIG 1B




-3/4-

FIG 1C



-4/4-

FIG 1D



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

**BLACK BORDERS**

**IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

**FADED TEXT OR DRAWING**

**BLURRED OR ILLEGIBLE TEXT OR DRAWING**

**SKEWED/SLANTED IMAGES**

**COLOR OR BLACK AND WHITE PHOTOGRAPHS**

**GRAY SCALE DOCUMENTS**

**LINES OR MARKS ON ORIGINAL DOCUMENT**

**REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

**OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**